


INFORMATION REPORTS

NUMBER 2010-3



FISH DIVISION

Oregon Department of Fish and Wildlife

Habrate: A Limiting Factors Model for Assessing Stream Habitat Quality
for Salmon and Steelhead in the Deschutes River Basin

Oregon Department of Fish and Wildlife prohibits discrimination in all of its programs and services on the basis of race, color, national origin, age, sex or disability. If you believe that you have been discriminated against as described above in any program, activity, or facility, or if you desire further information, please contact ADA Coordinator, Oregon Department of Fish and Wildlife, 3406 Cherry Drive NE, Salem, OR, 503-947-6000.

This material will be furnished in alternate format for people with disabilities if needed. Please call 541-757-4263 to request

Habrate: A Limiting Factors Model for Assessing Stream Habitat Quality for Salmon and Steelhead in the Deschutes River Basin

Jennifer L. Burke, Kim K. Jones, and Jeffrey M. Dambacher

Oregon Department of Fish and Wildlife
Corvallis, OR



Citation: Burke, J. L, K. K. Jones, and J. M. Dambacher. 2010. Habrate: A Limiting Factors Model for Assessing Stream Habitat Quality for Salmon and Steelhead in the Deschutes River Basin. Information Report 2010-03, Oregon Department of Fish and Wildlife, Corvallis.

CONTENTS

ABSTRACT	1
INTRODUCTION.....	2
STUDY AREA.....	5
ANADROMOUS SALMONIDS IN DESCHUTES RIVER BASIN	6
Historical distribution.....	6
METHODOLGY.....	7
Literature review and habitat criteria	7
Spreadsheet Components.....	7
Habitat Data.....	7
Rating Habitat.....	9
Input Criteria	12
Spatially-Explicite Output.....	13
RESULTS.....	13
Spatial Display of Reach Level Ratings.....	15
DISCUSSION	20
REFERENCES.....	22
Appendix A	32
Appendix B.....	33
Appendix C.....	40
Appendix D	46
Appendix E.....	48
Appendix F.....	51
Appendix G	54

FIGURES

Figure 1. Historic distribution (thin dark lines) and proposed reintroduction areas (thick dark lines) of anadromous fish in the mid- and upper Deschutes River basin. Significant passage barriers are labeled and depicted by filled circles.	3
Figure 2. Survey reaches (light gray lines) in the reintroduction area of the mid- and upper Deschutes River basin.....	8
Figure 3. Schematic of the HabRate assessment process and its elements.	10
Figure 4. Cumulative summary of reach ratings in seven subbasins in the Deschutes River. Each reach was given a rating of 1, 2, or 3. The count represents the number of reaches of a given score in each subbasin. Larger subbasins usually had more reaches. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.....	15
Figure 5. Reach level ratings of spawning, emergence and incubation habitat for steelhead, Chinook (top), steelhead (middle), and sockeye (bottom) in the Deschutes River basin above Lake Billy Chinook. Sockeye are only present in the Metolius drainage.	17
Figure 6. Reach level ratings of summer rearing habitat for chinook (top) and subyearling steelhead (bottom) in the Deschutes River basin above Lake Billy Chinook.	18
Figure 7. Reach level ratings of winter rearing habitat for chinook (top) steelhead (bottom) in the Deschutes River basin above Lake Billy Chinook.	19

TABLES

Table 1. Early life histories evaluated in HabRate.	6
Table 2. Reach attributes (averaged values) included in HabData	9
Table 3. Level 1 and 2 Reach attributes evaluated for Chinook and sockeye salmon and steelhead trout. *Excludes sockeye salmon. **Excludes steelhead trout.....	11
Table 4. Sample of input criteria section of worksheet for spawning, egg survival, and emergence of Chinook salmon. Non-shaded cells are adjustable and formulas throughout the worksheet update automatically through linked formulas.....	12
Table 5. Average HabRate ratings summarized by subbasins within the Deschutes River basin. Sockeye salmon spawn only in the Metolius basin, and rear in Suttle Lake and Lake Billy Chinook, which were not evaluated in HabRate. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.	14
Table 6. Sample of tiered results for Metolius River reaches evaluated for summer rearing life stage of subyearling steelhead trout. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.....	15
Table 7. Strengths and limitations of HabRate, a spatially explicit physical and biological limiting factors model.	21

ABSTRACT

Fishery managers are commonly tasked with the basic question “Will the contemporary habitat above a barrier support the fish populations that historically resided in the watershed?” Managers in central Oregon were confronted with that question in an effort to reestablish fish populations in 375 kilometers of stream above the Round Butte-Pelton Dam complex (Rkm 161) on the Deschutes River. Stream surveys had been conducted in most of the available stream habitat, but had not been synthesized in a form that allowed managers to view the quality and complexity of stream habitat in an easily-understandable fashion. In response, we developed a limiting factors model (HabRate) that assessed the potential quality of stream habitat using stream survey data for each juvenile life stage of salmon and steelhead. The model was developed for a specific application to the middle Deschutes River basin in Oregon, but was intended for general application to Pacific Northwest basins. To parametrize the model, we summarized available literature on salmonid habitat requirements. Habitat criteria were developed for discrete life history stages (i.e. spawning, egg survival, emergence, summer rearing, and winter rearing) and used to rate the quality of stream reaches as poor, fair, or good, based on attributes relating to stream substrate, habitat unit type, cover, gradient, temperature, and flow. Reach level summaries of stream habitat data were entered into MS Excel, and interpreted by a series of algorithms to provide a limiting factor assessment of potential egg-to-fry and fry-to-parr survival for each reach. Model output lists habitat quality by species and life stage for each reach of stream. The model is a decision making tool that is intended to provide a qualitative assessment of the habitat potential of stream reaches within a basin context. Design criteria for the model were simplicity, flexibility, and transparency. While HabRate was based on our interpretations of the published literature, specific criteria for habitat quality were structured to be easily adjusted where interpretations differ from ours. Information not common to standard stream survey designs, such as seasonal flow or temperature extremes can be included as input from professional judgment. The results were integrated into a GIS coverage coupled with the stream network and habitat data to provide a comprehensive map-based perspective of habitat quality in a watershed.

INTRODUCTION

The Deschutes River basin, located on the east flank of the north Oregon Cascades, formerly supported anadromous salmon and steelhead trout populations throughout the middle and upper reaches and tributaries. Salmon and trout populations first declined coincident with the degradation of river conditions and fish harvest pressure in the late 1800s. Deschutes River salmon and trout were subject to intense fishery harvest from the terminal fisheries on the lower Columbia River, and streams were dammed for irrigation withdrawal which blocked access to spawning and rearing habitat. Further habitat degradation resulted from the early transport of logs and associated activities along the Deschutes waterways. Construction of dams on the mainstem and tributary systems further restricted or reduced access to spawning and rearing areas. The construction of two small dams on Blue and Suttle lakes for recreational swimming extirpated the sockeye salmon populations in the lakes and in the Metolius River. The Crooked River basin, comprising two-thirds of the Deschutes River basin accessible to anadromous fish, was rendered inaccessible after the construction of Ochoco Dam in 1921 and Bowman Dam in 1961. Bonneville and Dalles dams on the mainstem Columbia River downstream of the Deschutes confluence, constructed in 1938 and 1960 respectively, decreased survival of outmigrant juveniles (Lichatowich et al. 1996). The final blow to anadromous fish in the Deschutes basin was the construction of the Pelton-Round Butte hydropower dam complex at RM 161 on the mainstem Deschutes River in 1958. Initial attempts to facilitate trout and salmon passage over the dams with fish ladders failed, leading to the removal of the ladder in 1968, a year marking the extinction of middle and upper Deschutes River salmon and trout populations (Nehlsen 1995).

In 1996, Portland General Electric (PGE) initiated the process to re-license the Pelton-Round Butte hydropower complex to continue operation in accordance with the Federal Energy Regulatory Commission guidelines (FERC). The application process required a plan for Protection, Mitigation and Enhancement Measures of environmental resources, particularly cultural resources and threatened and endangered species impacted by the complex. To comply with the license application guidelines, PGE committed to the reintroduction and establishment of Chinook and sockeye salmon and steelhead trout populations above the Pelton - Round Butte complex (Figure 1).




Figure 1. Historic distribution (thin dark lines) and proposed reintroduction areas (thick dark lines) of anadromous fish in the mid- and upper Deschutes River basin. Significant passage barriers are labeled and depicted by filled circles.

The reintroduction area covered a portion of the historic range of anadromous salmonids within the Crooked River, and all of the historic range within the Deschutes and Metolius Rivers (Figure 1). A feasibility study reviewed a number of biological and physical factors that may prevent the reestablishment of salmonids above the Round Butte-Pelton complex (Oosterhout 1999). One critical need was an evaluation of the suitability of aquatic habitat in each subbasin for each species and life stage of Chinook and sockeye salmon, and steelhead trout. The goal of the evaluation was to determine the likelihood that current aquatic conditions could support self-sustaining populations of anadromous salmonids, and if not, how and where restoration should occur. Our challenge, therefore, was to develop a

comprehensive and spatially explicit view of aquatic habitat conditions relevant to salmonid life history requirements for the three species.

In light of the decline of salmon populations in the Northwest and elsewhere, fish biologists have attempted to quantify the relationship of salmon life history and ecology with measurable attributes of aquatic habitat to predict productivity and survival. The ecological responses to physical conditions occurs at multiple scales, from micro to landscape scale (e.g. Vannote 1980; Hicks et al. 1991; Rahel and Hubert 1991; Nelson et al. 1992; Murray and Bailey 1998).

Seasonal use of specific habitat types by juvenile coho salmon has been used to predict potential carrying capacity of coastal streams for juvenile coho salmon in Oregon (Nickelson et al. 1992; Nickelson et al. 1993). Nickelson et al. (1992), Nickelson et al. (1993) and Nickelson and Lawson (1998) used channel habitat unit and reach level data coupled with temporal data (seasonal habitat use) to predict production and capacity at a stream and basin level. The coho salmon study was expanded to predict the viability of coho populations in three coastal basins in Oregon (Nickelson and Lawson 1998). However, a spatial or network component was not incorporated into the model.

Incorporating a life history approach to the relationship between habitat and biological response at a scale from channel habitat unit to river network adds another dimension to the interpretation (Lichatowich et al. 1995). The overall quality, connectivity, and relationships among habitats are crucial to the successful completion of a fish's life cycle. Modeling the survival of fish through the life cycle requires integrating spatial and temporal information. Kocik and Ferreri's (1998a) simulation model has been used to describe how the spatial structure of spawning and rearing habitat in a river system influenced the population dynamics of Atlantic salmon. Moberg et al. (1997) developed a spatially and temporally descriptive numerical model of the productivity and capacity of a system by integrating survival of a salmonid species at each life stage. The model then used potential life history trajectories of a salmonid to define connectivity within a watershed. A life history perspective has the advantage of incorporating spatial structure and connectivity of the habitat with the survival of fish at each life stage.

However, models that predict fish standing crop and production based on habitat parameters implicitly assume a deterministic relationship between fish and their physical environment. Such models are typically based either on regression analyses, or a limiting factors approach (Shrivell 1989). While some regression-based models have been highly predictive ($R^2 = 50$ to 96%) in the areas from which they were developed, their generality appears limited ($R^2 < 30\%$) when applied elsewhere without recalibration. Limiting factor models are applied with the implicit assumption that included variables are of general importance. Where the status of a particular population is poorly predicted, it is implied that the population is limited by variables not included in the model.

With the widespread application of the Hankin-Reeves stream survey design (Hankin and Reeves 1988), significant effort has been dedicated toward basin-wide assessments of stream habitat. Specifically in Oregon, an extensive stream survey program by the Oregon

Department of Fish and Wildlife (ODFW), Aquatic Inventories Project (AIP), has inventories of over 16,000 km of streams statewide (<http://oregonstate.edu/dept/ODFW/freshwater/inventory/basinwid.html>). The challenge remains to interpret an increasingly large volume of stream survey data in a way that is meaningful for basin-wide management of salmonid populations. While AIP stream survey data has been used to describe bull trout *Salvelinus confluentus* rearing areas (Dambacher and Jones 1994), and predict carrying capacity of juvenile coho salmon (Nickelson and Lawson 1998, Jones and Moore 1999), applications for other salmonid species have yet to be developed or researched. We sought therefore to derive meaningful criteria from existing literature of spawning and rearing habitat conditions based on life history studies for steelhead trout, chinook, and sockeye salmon. This effort grew out of a specific request to provide habitat-based stream production potential as input to a stochastic simulation of chinook and sockeye life history model for the Deschutes River basin (Oosterhooft 1999); however, our design is generally applicable to Pacific Northwest systems.

The model is a habitat rating system (HabRate) of aquatic conditions designed to link salmon with their environment and to provide a foundation for reach and watershed scale restoration programs. The conceptual basis for our modeling approach was to 1) capitalize on the wealth of existing field survey data (specifically basin survey data from ODFW), 2) describe habitat attributes at multiple scales, 3) be spatially explicit, 4) describe connectivity within a drainage, and, 5) build transparency and ease-of-use in the model to allow the user to adjust parameters and logic statements. HabRate describes habitat quality relative to each life stage of a salmonid species rather than numerically predict the carrying capacity of a habitat unit, reach, or stream. HabRate permits integration of survey and landscape data into a GIS format to display aquatic habitat within a watershed context and has incorporated flexibility of scale for comparisons between the reach, river, and basin level.

STUDY AREA

The Deschutes River basin lies adjacent to a major climate transition zone and within a geologically active landscape shaped by volcanism, tectonics, and glacial activity (Taylor and Hannan 1999). The Deschutes River system drains from the Eastern Cascade Mountain ecoregion in the west, the Blue Mountain ecoregion in the east, with the Northern Basin and Range ecoregion in the south and Columbia Plateau ecoregion in the north (Thorsen et al. 2003). The Cascades mountain range is volcanic in origin and historically contributed substantial amounts of lava and ash-tuff to the basin. The Blue Mountain Province is more open and lower elevation (Thorsen et al 2003), but is also primarily of volcanic origin. The river basin in the Cascades and southern region of the Blue Mountain ecoregion (John Day/Clarno lowlands and uplands) are unique in that the river primarily flows through lava fields pocketed by prairies. The Eastern Cascade and Blue Mountain Ecoregions receive a considerable amount of snow (greater than 80 inches per year on average) from November to March, while the remainder of the year is predominantly dry (Taylor and Hannan 1999). These regions support temperate alpine forests and meadows. The Northern Basin and Range and Columbia Plateau are largely composed of basalt and ash-tuff that have been eroded over time. In the Deschutes River Valley of the Columbia Plateau, the rivers flow through imposing basaltic canyons and arid meadows. The southern Blue Mountain and

Deschutes-Columbia Plateau are cool desert and steppe lands that receive less than 15 inches per year of precipitation in the lower elevations (Thorsen et al 2003, Taylor and Hannan 1999). The unique nature of the geology and climate of the region maintains river flows throughout the year. River flow throughout the Blue Mountain and Deschutes-Columbia Plateau is maintained by recurrent, and sometimes large, cold springs originating from snowmelt and precipitation percolating through a vast network of permeable volcanic rock.

ANADROMOUS SALMONIDS IN DESCHUTES RIVER BASIN

Historical distribution

HabRate describes the quality of habitat in the streams that were historically occupied by each life history stage of spring and fall Chinook salmon, sockeye salmon, and summer steelhead trout in the Deschutes River above the Pelton-Round Butte complex. The historical extent of chinook salmon lacked full documentation, although we pieced together the probable location and timing of adult and juvenile migration in the middle Deschutes basin (Appendix A). It is believed that spring Chinook spawned throughout the basin and summer/fall Chinook run spawned in the mainstem Deschutes and lower Metolius River (Nehlsen 1995) Progeny of the adult Chinook salmon runs were comprised of subyearling (ocean-type) and yearling (stream-type) migrants based on timing of their migration past the Pelton-Round Butte site (Nehlsen 1995). The spawning distribution of steelhead was poorly documented in the upper Deschutes basin, although they were thought to spawn throughout the accessible portions of the basin. The steelhead juveniles typically remained in the Deschutes River basin for 1 to 2 years (King 1966, Nelsen 1995). Sockeye salmon were confined to the Metolius River drainage, with probable spawning areas in Lake Creek and rearing in Suttle Lake.

Three life stages of Chinook and sockeye salmon and steelhead trout were evaluated; 1) spawning, incubation and emergence, 2) summer rearing, and 3) winter rearing. Spawning, incubation, and emergence were combined into a single life stage in the evaluation due to the similar criteria values (Table 1). Migratory conditions for adult salmonids were considered optimal for temperature and river flow; therefore, adult life history attributes were not evaluated. However, temperature and flow information can be incorporated in the HabRate model if available.

Table 1. Early life histories evaluated in HabRate.

Life History	Chinook Salmon	Steelhead Trout	Sockeye Salmon
Spawning, incubation, and emergence	X	X	X
Subyearling (0+) summer rearing	X	X	X
Subyearling (0+) overwintering	X	X	X
Yearling (1+) summer rearing		X	
Yearling (1+) overwintering		X	

METHODOLOGY

Literature review and habitat criteria

We performed an extensive literature review and compiled the habitat requirements of Chinook and sockeye salmon, and steelhead trout for each freshwater life history stage. Few juvenile salmonid life history studies were conducted in the Deschutes River basin. Consequently, the scope of the literature review for criteria values was expanded to include Alaska, Idaho, and the eastern regions of Oregon and Washington as necessary. We preferentially selected research from field studies over research in a laboratory setting. We evaluated three life history stages for chinook salmon. Spawning and 0+ summer rearing (limited duration) evaluation applied to both ocean-type and stream-type juveniles, while 0+ overwintering applied only to yearling (stream-type) juveniles. Five life stages of steelhead trout were evaluated that accounted for 1 to 2 years of freshwater rearing. Sockeye salmon had an abbreviated life history evaluation in HabRate, limited to spawning areas in streams with access to the expanded rearing potential in lakes.

The literature review is summarized by species: Chinook salmon (Appendix B), Steelhead trout (Appendix C), and sockeye salmon (Appendix D). Similarly, from these sources, we developed habitat rating criteria for each species, presented in Appendices E, F, and G, representing our interpretation of the various values presented in the literature.

Spreadsheet Components

HabRate is organized by four worksheet components; 1) HabData, 2) Evaluation, 3) Reach Rating, and 4) Input criteria. The elements that comprise each of the worksheets are discussed in the following sections. The model is available at <http://oregonstate.edu/dept/ODFW/freshwater/inventory/habratereg.htm>.

Habitat Data

Most of the streams and rivers in the Metolius, Deschutes and Crooked Rivers above the Round Butte complex available to anadromous salmonids have been surveyed (Figure 2). The analysis incorporated stream survey data from the Oregon Department of Fish and Wildlife (ODFW), Aquatic Inventories Project (Moore et al. 1997, 2007). Stream survey design followed the census approach described by Hankin (1984) and Hankin and Reeves (1988). While the primary objective of the Hankin (1984) and Hankin and Reeves (1988) methodology was to estimate the number of fish in a stream, it was adapted as a census survey design to efficiently collect information on aquatic habitat attributes continuously in a consistent format from the stream mouth to headwaters. This census survey design, frequently referred to as a basin survey, was a departure from the traditional representative reach survey for a basin (Dolloff et al. 1997). The major advantage to census surveys was the concurrent and continuous record of geomorphic reaches, habitat units, and associated features. It provided information on stream size, channel structure, large wood debris, sediment throughout the watershed, all features that influence the distribution and productivity of anadromous, fluvial, and resident fishes.

Habitat surveys in the middle Deschutes River basin began in 1989 by the USFS, and in 1993 by ODFW. Both agencies continued surveying in the basin through 1997, when HabRate was developed. Both agencies conducted the stream surveys during summer flow levels. Since 1997, ODFW has continued to survey streams, and these data have been incorporated into HabRate, replacing the older data as streams were resurveyed.




Figure 2. Survey reaches (light gray lines) in the reintroduction area of the mid- and upper Deschutes River basin.

Basin level stream surveys were completed in the Deschutes River from Lake Billy Chinook reservoir to Big Falls, the uppermost historic natural barrier. The Crooked River was surveyed from Lake Billy Chinook to Bowman Dam, and from the confluence with the Crooked River up Ochoco Creek to Ochoco Dam. Whychus Creek (Deschutes) and McKay Creek (Crooked) were also surveyed. In the Metolius watershed, the upper mainstem was surveyed as were most tributaries (Figure 2).

Reach level values consisted of total counts per reach, proportions (as a percentage) of the reach, averages, and counts per fixed length (100 meters or 1 km). The length of the reach was variable based on the geomorphology of the stream. The reach level values were compiled in MS Access and exported to a MS Excel worksheet, titled HabData on the worksheet tab, and served as the source data for the evaluation. The structure of HabRate is

such that the scope of the evaluation is expandable at any time to include additional reaches or streams by adding in additional rows of data.

HabData included 4 classes of data: substrate, channel morphology, habitat unit features, and large woody debris. Individual attributes within each category are listed in Table 2. All of the attributes were compiled in HabData in metric units. Although not all attributes were used in the analysis, we retained them for reference. The habitat rating process evaluated stream habitat attributes collectively deemed important for productivity at each life stage (Table 3).

Table 2. Reach attributes (averaged values) included in HabData.

Substrate	Channel Morphology	Habitat	Wood
Percent fines	Reach length	Number of pools	Pieces of large woody debris (LWD)
Percent gravel	Channel area	Percent pools	Volume of LWD
Percent cobble	Gradient	Scour pool depth	Pieces of LWD per 100m
Percent boulders	Wetted width	Depth of riffles	Volume of LWD per 100m
Percent fines in riffles	Active channel width	Pools per km	Key pieces of LWD
Percent gravel in riffles	Large boulders	Pools greater than 1m depth per km	Key pieces of LWD per 100m
Average percent boulders per pool	Large boulders per 100m	Channel width (bankfull) pools	Average LWD per pool
	Percent Open sky	Number of Pools per 100m	Average key pieces of LWD per pool
	Width to depth ratio	Residual pool depth	
		Percent undercut	
		Average percent undercut per pool	

Rating Habitat

HabRate hierarchically rates and then evaluates the attributes at three levels (Figure 3, Table 3). At Level 1, a rating is generated for each individual attribute for each applicable life history of each species (Table 3). In Level 2, the rating summarized the attributes by category to represent the collective condition. In the final Level 3, the rating evaluates Level 2 rating values using a combination of individual and collective assessments. For instance, spawning, incubation, and emergence rating evaluates the substrate separately from the combined evaluation of pool area and residual pool depth ratings. The approach focuses the rating on conditions potentially inadequate in quality and survival, without compromising the value of equally important habitat features for an overall rating. The model retains each subcomponent for reference.




Figure 3. Schematic of the HabRate assessment process and its elements.

Table 3. Level 1 and 2 Reach attributes evaluated for Chinook and sockeye salmon and steelhead trout. *Excludes sockeye salmon. **Excludes steelhead trout.

Level 1		Level 2	
Spawning, Incubation, Emergence	Percent fines	Substrate	
	Percent gravel		
	Percent cobble		
	Pool area*	Morphology	
	Residual pool depth		
	Gradient**		
Summer Rearing (Chinook salmon and steelhead trout only)	Percent fines	Substrate	
	Percent gravel**		
	Percent cobble and boulders		
	Pool area	Pool area	
	Average scour pool depth per pool**	Pool complexity**	Pool Complexity**
	Average large woody debris per pool**		
	Undercut	Cover	
	Large woody debris per 100m		
	Large boulders per 100m		
	Gradient**	Gradient	
	Overwintering (Chinook salmon and steelhead trout only)	Percent fines	Interstices
Percent cobble and boulders			
Pool area		Pool habitat	
Average scour pool depth per pool**			Pool complexity**
Average large woody debris per pool**			
Percent undercut		Cover	
Large woody debris per 100m			
Large boulders per 100m			
Gradient		Gradient	
Summer rearing 1+ (steelhead trout only)	Percent fines	Interstices	
	Percent cobble and boulders	Pool Habitat	
	Pool area		
	Depth in fast water units	Hydrology	
	Percent undercut	Cover	
	Large woody debris per 100m		
	Large boulders per 100m		
Overwintering 1+ (steelhead trout only)	Percent fines	Interstices	
	Percent cobble and boulders	Pool Habitat	
	Pool area		
	Percent undercut	Cover	
	Large woody debris per 100m		
	Large boulders per 100m		

Input Criteria

Each Level attribute is evaluated using a range of criteria and assigned a numerical value on a scale of 1 to 3, with 3 being the best condition. A rating of 1 equated to poor conditions (potential low survival) that are very detrimental to the eggs or juveniles. A rating of 2 reflected conditions favorable (fair) to survival or adequate for juvenile use. A rating of 3 indicated conditions were optimum for productivity or survival.

We constructed a criteria input table for each life history stage of each species that show the criteria and logic statements used to rate individual attributes for Level 1 (Table 4). The structure of the criteria input worksheet allows for easy adjustments of the critical values, denoted by white boxes, which link the criteria input worksheet to the HabData worksheet and rating worksheets. This permits the user to adjust the criteria ranges, if deemed necessary, which will automatically update the formulas and adjusts the resultant ratings in the subsequent rating worksheet. As structured, the rating process has a greater geographic range of application through the adjustment and refinement of criteria values according to ecological province. Please note the input page was not intended for use in a 'what if' scenario, which could lead to erroneous interpretations and results.

Table 4. Sample of input criteria section of worksheet for spawning, egg survival, and emergence of Chinook salmon. Non-shaded cells are adjustable and formulas throughout the worksheet update automatically through linked formulas.

Attribute	Criteria and Rating		
	3	2	1
Fines (%)	≤ <input type="text" value="10"/>	> 10 and ≤ 20	> <input type="text" value="20"/>
Gravel (%)	≥ <input type="text" value="30"/>	< 30 and > 15	≤ <input type="text" value="15"/>
Cobble (%)	≥ <input type="text" value="20"/> and ≤ <input type="text" value="40"/>	< 20 and ≥ 10 > 40 and ≤ 70	< <input type="text" value="10"/> or > <input type="text" value="70"/>
Pool Area (% pools)	≥ <input type="text" value="40"/> and ≤ <input type="text" value="60"/>	< 40 and ≥ 20	< <input type="text" value="20"/> or > 60
Residual Pool depth (m)	≥ <input type="text" value="0.2"/>		< 0.2
Gradient (%)	< <input type="text" value="4"/>		≥ 4

The scaled range (e.g. 1, 2, or 3) and minimum value rating methodology was preferred to mean values so as not to obscure potentially detrimental attributes, and to identify limiting factors in each reach. Because the rating process is adjustable and transparent, the analysis can identify individual attributes responsible for a low rating in a given reach for a species' life stage.

All habitat surveys were conducted during summer flows, levels that are typically the lowest for the Deschutes River basin. In the evaluation of the winter habitat component of chinook salmon and steelhead trout, the depth criteria was interpreted from winter flow studies. The summer data were extrapolated to represent base flow during winter conditions.

Temperature and flow conditions were included in the list of attributes. Due to the nature of the Deschutes River basin, winter temperature and discharge were not considered a limiting factor for early life history development. Very little data existed evaluating the effects of summer temperatures and flows on juvenile salmonids in the Deschutes River. Therefore, those variables were not included in the evaluation although the formulas in HabRate retained the variables for use in other basins

where irrigation withdrawals or natural fluctuations in flow may create critical thresholds for flow or temperature.

Spatially-Explicit Output

HabRate is structured to link the components and the final output to a spatially-explicit GIS dataset to provide the rating results in a map-based view. The structure and model results integrate with a geospatial stream reach dataset using a unique identifier for each reach, i.e. LLID code and reach number (HABRCH). The habitat quality rating and habitat data for each reach, species, and life history can then be displayed in a Geographic Information System (GIS) on a digitized stream layer. Even though connectivity is not modeled between reaches, the map-based view of ratings and attributes permits additional analysis of the spatial connectivity and salmonid survival between reaches.

RESULTS

HabRate generated a rating of salmonid habitat quality and potential limiting factors in the Deschutes River basin for three species of salmonids and one to five life histories per species. While the focus was on the reintroduction area in the mid- and upper Deschutes basin, we also evaluated the streams in the lower Deschutes to provide a balance and perspective on conditions throughout the Deschutes basin. The average reach level rating per species and subbasin provide an initial assessment of average conditions of the surveyed reaches for each life history (Table 5). Ratings that are less than 2 were subbasins with substandard conditions for that particular life history. For example, Buckhollow Creek basin rated the lowest for spawning, incubation, and emergence for steelhead trout, while the remaining steelhead life histories rated predominately adequate to optimum. Here we will provide selected examples of how the model functions and selected examples of the output.

Figure 4 displays the cumulative counts of reach ratings (1, 2, or 3) for the spawning, incubation, and emergence life stage for chinook salmon in seven subbasins in the Deschutes River. For example the scores from 14 reaches in Bakeoven subbasin were summarized to display the frequency of scores for that life stage. Most of the habitat was good for spawning, incubation, and emergence. Table 6 provides an example of the level of detail that goes into each reach rating; in this case for the summer rearing of subyearling steelhead trout in the Metolius River subbasin. The survey variables were summarized at level 1, then combined by logic statements to level 2, and finally combined through logic statements to an overall rating for the reach for that life stage and species. The approach allows the rating to be dissected to their component indices or variables to gain a better understanding of the quality of habitat. In the same way, the component ratings, such as large wood or fine sediment, could be displayed in GIS to demonstrate areas of limited complexity or excessive sedimentation.

Table 5. Average HabRate ratings summarized by subbasins within the Deschutes River basin. Sockeye salmon spawn only in the Metolius basin, and rear in Suttle Lake and Lake Billy Chinook, which were not evaluated in HabRate. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.

Species	Spawn and Emergence	Summer Rearing	Overwintering	Summer Rearing 1+	Overwinter 1+
<i>Bakeoven Creek</i>					
Steelhead salmon	2.8	2.3	2.9	2.0	2.9
Chinook salmon	2.7	2.0	2.4	-	-
Sockeye salmon	-	-	-	-	-
<i>Buckhollow Creek</i>					
Steelhead salmon	1.3	2.3	2.5	1.8	2.5
Chinook salmon	2.3	2.0	2.2	-	-
Sockeye salmon	-	-	-	-	-
<i>Crooked River</i>					
Steelhead salmon	1.6	2.3	2.3	1.9	2.4
Chinook salmon	1.9	2.0	2.2	-	-
Sockeye salmon	-	-	-	-	-
<i>Metolius River</i>					
Steelhead salmon	1.6	2.2	2.1	2.1	2.4
Chinook salmon	1.6	1.6	1.9	-	-
Sockeye salmon	1.8	-	-	-	-
<i>Shitike Creek</i>					
Steelhead salmon	1.8	2.6	2.5	1.9	2.6
Chinook salmon	1.8	1.9	2.0	-	-
Sockeye salmon	-	-	-	-	-
<i>Trout Creek</i>					
Steelhead salmon	2.1	2.4	2.5	1.9	2.7
Chinook salmon	2.3	2.0	2.0	-	-
Sockeye salmon	-	-	-	-	-
<i>Upper Deschutes River</i>					
Steelhead salmon	1.9	2.2	2.6	2.0	2.6
Chinook salmon	2.2	2.0	2.4	-	-
Sockeye salmon	-	-	-	-	-
<i>Warm Springs River</i>					
Steelhead salmon	1.9	2.6	2.5	2.0	2.7
Chinook salmon	2.0	2.0	2.1	-	-
Sockeye salmon	-	-	-	-	-




Figure 4. Cumulative summary of reach ratings in seven subbasins in the Deschutes River. Each reach was given a rating of 1, 2, or 3. The count represents the number of reaches of a given score in each subbasin. Larger subbasins usually had more reaches. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.

Table 6. Sample of tiered results for Metolius River reaches evaluated for summer rearing life stage of subyearling steelhead trout. A value of 1 = poor, 2 = fair, and 3 = good habitat quality.

Stream	Reach	Level 1					Level 2			Rating
		Fines	Cobble and boulders	Undercut	Large woody debris per 100m	Boulders per 100m	Substrate	Pool Area	Cover	
Mariel Creek	1	2	2	2	3	1	2	1	2	2
Mariel Creek	2	3	3	1	2	3	3	1	2	2
Mariel Creek	3	1	2	1	1	1	2	1	1	2
Metolius River	1	2	3	3	1	1	3	2	2	3
Metolius River	2	2	2	3	1	1	2	1	2	2
Parker Creek	1	3	3	3	3	2	3	1	3	3
Parker Creek	2	3	3	1	1	3	3	1	2	2
Whitewater River	1	2	3	1	2	3	3	1	2	2
Whitewater River	2	1	3	1	3	3	2	1	3	2
Whitewater River	3	1	3	1	3	3	2	1	3	2

Spatial Display of Reach Level Ratings

Because the HabRate results are referenced to spatially-explicit hydrologic datasets, the results may be mapped at all levels. Reach level ratings provide the coarsest resolution to assess the spatial distribution of the HabRate evaluation (e.g. Figure 7). Individual metrics (e.g. complex pools, high quality spawning habitat) could also be mapped at a channel unit scale, on the order of tens of meters rather than kilometers. Figures 7, 8, and 9 display the distribution of habitat quality relative to spawning and emergence, summer rearing, and

winter rearing, respectively. The variability in habitat quality is apparent within and between each subbasin. For example, the mainstem Crooked River has low quality habitat for spawning and emergence, and moderate quality for rearing through much of its length. However, McKay Creek, a tributary to the Crooked River has fair and good quality habitat for all life stages.




Figure 5. Reach level ratings of spawning, emergence and incubation habitat for steelhead, Chinook (top), steelhead (middle), and sockeye (bottom) in the Deschutes River basin above Lake Billy Chinook. Sockeye are only present in the Metolius drainage.




Figure 6. Reach level ratings of summer rearing habitat for chinook (top) and subyearling steelhead (bottom) in the Deschutes River basin above Lake Billy Chinook.




Figure 7. Reach level ratings of winter rearing habitat for chinook (top) steelhead (bottom) in the Deschutes River basin above Lake Billy Chinook.

DISCUSSION

HabRate, developed in 1997, is one of many models to evaluate suitability of salmonid habitat as part of conservation, restoration, and reintroduction efforts. Here we presented a model that permitted an examination of habitat features that influence productivity and capacity for different life history strategies and life stages of salmon and steelhead in the Deschutes basin. We built this model based on the availability of an extensive and spatial explicit data set of stream habitat conditions in the study area coupled with literature rich with information on habitat requirements of salmonids. Due to the limited availability of field-collected stream data in most areas, many salmonid habitat assessments employ best professional judgment in the rating procedure, often at the landscape-scale or watershed-scale using decision-supported logic models that may or may not included field-collected data. Examples include the widely used Ecosystem Diagnosis and Treatment (EDT) developed by Moberg Biometrics Inc. (2003), and the Ecosystem Management Decision Support (EMDS) developed and used by Reynolds and Peets (2001) on the Chewaucan River basin in Southern Oregon. These habitat evaluation models are often not transparent or straightforward but are capable of complex assessments and analyses that may incorporate dozens of variables. This model, in contrast, was developed specifically to take advantage of the availability of documented field data, and to use literature-based relationships of habitat to fish.

Less complex, linear evaluations that utilize best professional judgment in the review of habitat criteria within a river basin are also widely used and closely resemble the structure of HabRate. An example includes Smith's (2005) Washington statewide stream-level assessment for the State of the Salmon report that included critical limiting factors ratings per stream and summarized at the watershed level.

Although HabRate is not novel in its objectives, it is unique in its compatibility with the Aquatic Inventories Project Stream Survey methods and results, transparency throughout the analysis, and direct link to GIS, including the original stream survey data at the unit, i.e. riffle, pool, etc., reach levels attribute data, and each rating level of the HabRate analysis. In addition, all HabRate criteria and logic statements are editable in a MS Excel spreadsheet for easy updates and application to basins outside of the Deschutes River. Furthermore, stream survey data can be continuously added and updated easily at any time, providing a unique monitoring tool that works readily with Microsoft Access, Microsoft Excel, and ArcGis. The model criteria can also readily updated as new research provides additional insight on the relationship of habitat attributes to survival at different life stages, and the model can be modified to provide habitat quality ratings for Chinook, steelhead, and sockeye in other provinces or adapted to other salmon species. Additional sources of information can be integrated or overlaid with the habitat data such as water quality (e.g. temperature), instream structures (e.g. passage barriers, diversion structures), or landscape features (e.g. geology, land use) to provide a more comprehensive perspective on the basin.

Different modeling methods inherently possess different strengths and limitations. We compiled a list identifying HabRate's strengths and limitations in Table 7.

Table 7. Strengths and limitations of HabRate, a spatially explicit physical and biological limiting factors model.

Strengths	Limitations
Uses quantitative and qualitative data	No modeled connectivity between reaches
Flexible scales	No empirical testing of results
Visual presentation (GIS)	No multiplicative effects or interactions
User adjustable criteria	Static evaluation (discreet to life history stage)
Wide geographic range of application	Single species evaluation
Life history stage breakdown	Limited by the quality of data available
Simple - straightforward evaluation	
Identifies potential limiting factors	
Spatial relationships	
Transparent evaluation process	

HabRate was developed as tool to evaluate the suitability of habitat for salmonids in the Deschutes River basin, but has a much broader application to other basins in the Pacific Northwest. Because HabRate provides a link between stream habitat conditions and life history requirements of salmon, it may be used as is or adapted to identify limiting factors in stream conditions for prioritizing habitat restoration and developing recovery plans for salmon and trout in other basins

REFERENCES

- Bell, M.C. 1986. Fisheries handbook of engineering requirements and biological criteria. Fish passage development and evaluation program. US Army Corps of Engineers, North Pacific Division. 290 pp.
- Benhke, R. J. 1992. Native trout of western North America. American Fisheries Society, Monograph 6, Bethesda, Maryland.
- Binns, N.A. and F.M. Eiserman. 1979. Quantification of fluvial trout habitat in Wyoming. Transactions of the American Fisheries Society 108(3): 215-228.
- Bisson, P.A., J.L. Nielsen, R.A. Palmason, and L.E. Grove. 1981. A system of naming habitat types in small streams, with examples of habitat utilization salmonids during low streamflow. In: Armantrout, N.B, editor. Acquisition and utilization of aquatic habitat inventory information. Bethesda, Maryland: American Fisheries Society. p 62-73.
- Bisson, P.A., K. Sullivan, and J.L. Nielsen. 1988. Channel hydraulics, habitat use, and body form of juvenile coho salmon, steelhead, and cutthroat trout in streams. Transactions of the American Fisheries Society 117: 262-273.
- Bjornn, T.C. 1968. Survival and emergence of trout and salmon fry in various gravel-sand mixtures. In: Logging and salmon,: proceedings of a forum. Juneau : American Institute of Fishery Biologists, Alaska District. pp. 80 - 88.
- Bjornn, T.C. 1969. Embryo survival and emergence studies. Job No. 5, Federal Aid in Fish and Wildlife Restoration. Job Completion Report, project F-49-R-7. Idaho Fish and Game Department, Boise. 11 pp.
- Bjornn, T.C. 1971. Trout and salmon movements in two Idaho streams as related to temperature, food, stream flow, cover, and population density. Transactions of the American Fisheries Society 100:423 - 438.
- Bjornn, T.C., M.A. Brusven, M.P. Molnau, J.H. Milligan, R.A. Klamt, E. Chaco, and C. Schaye. 1977. Transport of granitic sediment in streams and its effect on insects and fish. University of Idaho, Forest, Wildlife and Range Experiment Station Bulletin 17, Moscow.
- Bjornn, T.C., and D.W. Reiser. 1991. Habitat requirements of salmonids in streams. In: W.R. Meehan., Editor. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitat. Bethesda, Maryland. American Fisheries Society Special Publication 19: 83-138.
- Black, E.C. 1953. Upper lethal temperatures of some British Columbia freshwater fishes. Journal of the Fisheries Research Board of Canada 10:196-210.

- Brett, J.R. 1952. Temperature tolerance in young Pacific salmon, genus *Oncorhynchus*. Journal of the Fisheries Research Board of Canada 9:265-323.
- Brett, J.R., W.C. Clar, and J.E. Shelbourn. 1982. Experiments on the thermal requirements for growth and food conversion efficiency of juvenile chinook salmon. Canadian Technical Report of Fisheries and Aquatic Sciences 1127. Pacific Biological Station, Nanaimo, BC. 29 pp.
- Briggs, J.C. 1953. The behavior and reproduction of salmonid fishes in a small coastal stream. California Department of Fish and Game, Fisheries Bulletin 94. 62 pp.
- Brusven, M.A., W.R. Meehan, and J.F. Ward. 1986. Summer use of simulated undercut banks by juvenile chinook salmon in an artificial Idaho channel. North American Journal of Fisheries Management 6:32-37.
- Burger, C.V., D.B. Wangaard, R.L. Wilmot, and A.N. Palmisano. 1982. Salmon investigations in the Kenai River, Alaska, 1979-1981. USDI Fish and Wildlife Service, National Fisheries Research Center. Seattle, Washington. 178 pp.
- Burner, C.J. 1951. Characteristics of spawning nests in Columbia River salmon. U.S. Fish and Wildlife Service Bulletin 52:96-110.
- Bustard, D.R. and D.W. Narver. 1975. Aspects of the winter ecology of juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). Journal of Fisheries Research Board of Canada. 32: 667-680.
- Chambers, J.S., R.T. Pressey, J.R. Donaldson, and W.R. McKinley. 1954. Research relating to study of spawning grounds in natural areas. Annual Report, contract DA.35026-ENG-20572, Washington State Department of Fisheries, Olympia.
- Chambers, J.S., G.H. Allen, and R.T. Pressey. 1955. Research relating to study of spawning grounds in natural areas. Annual Report, contract DA.35026-ENG-20572, Washington State Department of Fisheries, Olympia.
- Chambers, J.S. 1956. Research relating to study of spawning grounds in natural areas. U.S. Army, Corps. of Engineers, North Pacific Division, Fisheries Engineering Research Program. pp. 88-94.
- Chapman, D.W. and T.C. Bjornn. 1969. Distribution of salmonids in streams with special reference to food and feeding. In: Northcote, T.G., editor. Symposium on salmon and trout in streams. Vancouver: H.R. MacMillan Lectures in Fisheries, University of British Columbia. pp 153-176.
- Divinin, P.A. 1952. The salmon of South Sakhalin. Izvestiia Tikhookeanskogo Nauchno-Issledovatelskogo. Instituta Rybnogo Khoziaistvo i Okeanografii. (Izv. TINRO), 37:69-108.

- Dolloff, C. A., H. E. Jennings, and M. D. Owen. 1997. A comparison of basinwide and representative reach habitat survey techniques in three southern Appalachian watersheds. *North American Journal of Fisheries Management* 17:339-347.
- Edmundson, E.F., F.H. Everest, and D.W. Chapman. 1968. Permanence of station in juvenile chinook salmon and steelhead trout. *Journal of the Fisheries Research Board of Canada* 25(7):1453-1464.
- Eddy, R.M. 1972. The influence of dissolved oxygen concentration and temperature on survival and growth of chinook salmon embryos and fry. MS thesis. Oregon State University, Corvallis. 45 pp.
- Everest, F.H. 1969. Habitat selection and spatial interaction of juvenile chinook salmon and steelhead trout in two Idaho streams. Ph.D. dissertation, University of Idaho, Moscow. 77 pp.
- Everest, F.H., and D.W. Chapman. 1972. Habitat selection and spatial interaction by juvenile chinook salmon and steelhead trout in Two Idaho streams. *Journal of Fisheries Research Board of Canada*. 29: 91-100.
- Everest, F.L., R. L. Beschta, J.C. Scrivener, K.V. Koski, J.R. Sedell, and C.J. Cederholm. 1987. Fine sediment and salmonid production - a paradox. In: Salo, E.O, and T.W. Cundy, editors. *Streamside management: forestry and fishery interactions*. Seattle: College of Forest Resources, University of Washington. pp. 98 - 142.
- Fausch, K.D. 1993. Experimental analysis of microhabitat selection by juvenile steelhead (*Onchorynchus mykiss*) and coho salmon (*O. kisutch*) in a British Columbia Stream. *Canadian Journal of Fisheries and Aquatic Sciences*. 50: 1198-1207.
- Frey, 1942. Field notes summarized in the report by Nielson 1950. Cited in Nehlsen 1995.
- Greeley, J.R. 1932. The spawning habits of brook, brown, and rainbow trout and the problem of egg predators. *Transactions of the American Fisheries Society* 62: 239-248.
- Groot, C. and L. Margolis. *Pacific Salmon: Life Histories*. 1991. UBC Press, British Columbia, Canada. 576 pp.
- Hall, J.D., and Lantz, R.L. 1969. Effects of logging on the habitat of coho salmon and cutthroat trout in coastal streams. In: Northcote, T.G., editor. *Symposium on salmon and trout in streams*. Vancouver: H.R. MacMillan Lectures in Fisheries, University of British Columbia. p. 355-375.
- Hanel, J. 1971. Official memo to Dr. J.A.R. Hamilton. Pacific Power and Light Co., Portland, Oregon. July 14, 1971. Subject: Iron Gate fish hatchery steelhead program. 20 pp.
- Hankin, D. G. 1984. Multistage sampling designs in fisheries: applications in small streams. *Canadian Journal of Fisheries and Aquatic Sciences* 41:1575-1591.

- Hankin, D. G., and G. H. Reeves. 1988. Estimating total fish abundance and total habitat area in small streams based on visual estimation methods. *Canadian Journal of Fisheries and Aquatic Sciences* 45:834-844.
- Hartman, W.L., C.W. Strickland, and D.T. Hoopes. 1962. Survival and behavior of sockeye salmon fry migrating into Brooks Lake, Alaska. *Transactions of the American Fisheries Society* 91(2): 133-141.
- Hartman, G. F. 1965. The role of behavior in the ecology and interaction of underyearling coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada* 22: 1035-1081.
- Hicks, B.J. 1990. The influence of geology and timber harvest on channel morphology and salmonid populations in Oregon Coast Range streams. Ph.D. dissertation. Oregon State University, Corvallis. p. 212.
- Hicks, B. J., J. D. Hall, P. A. Bisson, and J. R. Sedell. 1991a. Responses of salmonids to habitat changes. Pages 483-518. *In: Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*. Meehan, W. R. (ed.) American Fisheries Society. Bethesda, Maryland, USA.
- Hillman, T.W., and J.S. Griffith. 1987. Summer and winter habitat selection by juvenile chinook salmon in a highly sedimented Idaho stream. *Transactions of the American Fisheries Society* 116: 185-195.
- Hunter, J.W. 1973. A discussion of game fish in the state of Washington as related to water requirements. Report by the Washington State Game, Fishery Management Division to the Washington State Department of Ecology, Olympia. 66 p.
- Huntington, C.W. 1985. Deschutes river spawning gravel study, vol. 1 & 2. Contract No. DE-AC79-83BP13102. Buell & Associates, Beaverton, Oregon. 850 pp.
- Johnson, S.W., J. Heifetz, and K.V. Koski. 1986. Effects of logging on the abundance and seasonal distribution of juvenile steelhead in some southeastern Alaska streams. *North American Journal of Fisheries Management* 6:532-537.
- Johnson, S.W., J.F. Thedinga, and K.V. Koski. 1992. Life history of juvenile ocean-type chinook salmon (*Onchorynchus tshawytscha*) in the Situk River, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*. 49: 2621-2629.
- Jonasson, B.C., J.V. Tranquilli, M. Keefe, and R.W. Carmichael. 1995. Investigations into the early life history of naturally produced spring chinook salmon in the Grande Ronde River Basin. Oregon Department of Fish and Wildlife, Annual Progress Rept BPA Project Number 92-026-04.

- Jonasson, B.C., J.V. Tranquilli, M. Keefe, and R.W.Carmichael. 1996. Investigations into the early life history of naturally produced spring chinook salmon in the Grande Ronde River Basin. Oregon Department of Fish and Wildlife, Annual Progress Rept BPA Project Number 92-026-04.
- Jonasson, B.C., J.V. Tranquilli, M. Keefe, and R.W.Carmichael. 1997. investigations into the early life history of naturally produced spring chinook salmon in the Grande Ronde River Basin. Oregon Department of Fish and Wildlife, Annual Progress Rept BPA Project Number 92-026-04.
- Jonasson, B.C., J.V. Tranquilli, M. Keefe, and R.W.Carmichael. 1998. Investigations into the early life history of naturally produced spring chinook salmon in the Grande Ronde River Basin. Oregon Department of Fish and Wildlife, Annual Progress Rept BPA Project Number 92-026-04.
- Jones, K. K. and K. M. S. Moore. 1999. Habitat assessment in coastal basins in Oregon: implications for coho salmon production and habitat restoration. Pages 329-340 In Sustainable Fisheries Management. E.E. Knudsen, C.R. Steward, D.D. McDonald, J.E. Williams, and D.W. Riser Eds. CRC Press New York 724 p.
- King, D.N. 1966. Central Region Administrative Report No. 66-3. Deschutes River summer steelhead. March. Oregon State Game Commission. Portland, Oregon. Cited in Nehlsen 1995
- Kocik, J. F. and C. P. Ferreri. 1998. Juvenile production variation in salmonids: populations dynamics, habitat, and the role of spatial relationships. Canadian Journal of Fisheries and Aquatic Sciences 55(Supplement 1): 191-200.
- Kondolf, G.M., and M.G. Wolman. 1993. The sizes of salmonid spawning gravels. Water Resources Research. 29: 2275-2285.
- Kondolf, G.M., M.J. Sale, and M.G. Wolman. 1993. Modification of fluvial gravel size by spawning salmonids. Water Resources Research. 29: 2265-2274.
- Konopacky, R.C. 1984. Sedimentation and productivity in a salmonid stream. Doctoral dissertation. University of Idaho, Moscow.
- Lagler, K.F. 1956. Freshwater fishery biology. Dubuque, IA: Wm C. Brown Co. 421 pp.
- Lewis, S.L. 1969. Physical factors influencing fish populations in pools of a trout stream. Transactions of the American Fisheries Society. 98(1): 14-19.
- Lichatowich, J., Mobernd, L., Lestelle, L., and Vogel, T. 1995. An approach to the diagnosis and treatment of depleted Pacific salmon populations in Pacific Northwest watersheds. Fisheries 20(1):10-18.

- Lichatowich, J. A., L. E. Mobrand, R. J. Costello, and T. S. Vogel. 1996. A history of the frameworks used in the management of Columbia River chinook salmon. Bonneville Power Administration. Contract No. DE-AM79-92BP25105. 86 pp.
- Lister, D.B. and H.S. Genoe. 1970. Stream habitat utilization by cohabitating underyearlings of chinook (*Oncorhynchus tshawytscha*) and coho (*Oncorhynchus kisutch*) salmon in the Big Qualicum River, British Columbia. Journal of the Fisheries Research Board of Canada 27(7): 1215-1224.
- Lorenze, J.M., and J.H. Eiler. 1989. Spawning habitat and redd characteristics of sockeye salmon in the Glacial Taku River, British Columbia and Alaska. Transactions of the American Fisheries Journal. 118: 495-502.
- Lotspeich, F.B. and F.H. Everest. 1981. A new method for reporting and interpreting textural composition of spawning gravel. USDA Forest Service, Research Technical Note PNW-369, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon.
- Lucas, K.C. 1959. The Robertson Creek spawning channel. Canadian Fish Culturist 25:4-23.
- Lunetta, R.S., B.L. Cosentino, D.R. Montgomery, E.M. Beamer, and T.J. Beechie. 1997. GIS - based evaluation of salmon habitat in the Pacific Northwest. Photogrammetric Engineering and Remote Sensing. 63(10):1219-1229.
- Mattson, C.K. 1948. Spawning ground studies of Willamette River spring chinook salmon. Oregon Fish Commission Research Briefs 1(2):21-32.
- McCuddin, M.E. 1977. Survival of salmon and trout embryos and fry in gravel-sand mixtures. MS thesis, University of Idaho, Moscow. 30 pp.
- McDonald, J. 1960. The behavior of Pacific salmon fry during their downstream migration to freshwater and saltwater nursery areas. Journal of the Fisheries Research Board of Canada 17:655-676.
- McNeil, W.J. and W.H. Ahnell. 1964. Success of pink salmon spawning relative to size of spawning bed materials. U.S. Fish and Wildlife Service, Special Scientific Report of Fisheries, No. 469. 15p.
- Meehan, W.R., and T.C. Bjornn. 1991. Salmonid distributions and life histories. In: Meehan, W.H., editor. Influences of forest and rangeland management on salmonid fishes and their habitat. Bethesda: American Fisheries Society Special Publication 19. p. 47-82.
- Mobrand, L. E., J. A. Lichatowich, L. C. Lestelle, and T. S. Vogel. 1997. An approach to describing ecosystem performance "through the eyes of salmon." Can. J. Fish. Aquat. Sci. 54: 2964-2973.

- Mobrand, L.E. 2003. Ecosystem Diagnosis and Treatment. <http://www.mobrand.com/edt>.
- Moore, K. M. S., K. K. Jones, and J. M. Dambacher. 1997. Methods for stream habitat surveys. Oregon Department of Fish and Wildlife, Information Report 97-4, Portland, Oregon.
- Moore, K.M.S, K.K. Jones, and J.M. Dambacher. 2007. Methods for Stream Habitat Surveys: Aquatic Inventories Project. Information Report 2007-01, version 3, Oregon Department of Fish & Wildlife, Corvallis. 67p.
- Murray, C.B., and M.L. Rosenau. 1989. Rearing of juvenile chinook salmon in nonnatal tributaries of the Lower Fraser River, British Columbia. Transactions of the American Fisheries Society. 118: 284 - 289.
- Murray, G. R. and R. J. Bailey (eds.). 1998. Change in Pacific Northwest coastal ecosystems. Proceedings of the Pacific Northwest Ecosystems Regional Study Workshop, August 13-14, 1996, Troutdale, OR. NOAA Coastal Ocean Program Decision Analysis Series No. 11. NOAA Coastal Ocean Office, Silver Spring, MD.
- Nehlsen, W. 1995. Historic salmon and steelhead runs of the upper Deschutes River and their environments. 65 pp. Copies available from: PGE Co., Hydro Licensing Department, attn: Marty May, 3WTCBR04, 121 SW Salmon St., Portland, Oregon, 97204.
- Nelson, R. L., W. S. Platts, D. P. Larsen, and S. E. Jensen. 1992. Trout distribution and habitat in relation to geology and geomorphology in the North Fork Humboldt River drainage, northeastern Nevada. Trans. Am. Fish. Soc. 121:405-426.
- Nickelson, T. E. and P. W. Lawson. 1998. Population viability of coho salmon, *Oncorhynchus kisutch*, in Oregon coastal basins: application of a habitat-based life cycle model. Can. J. Fish. Aq. Sci. 55: 2383-2392.
- Nickelson, T. E., J. D. Rodgers, S. L. Johnson, and M. F. Solazzi. 1992. Seasonal changes in habitat use by juvenile coho salmon (*Oncorhynchus kisutch*) in Oregon coastal streams. Can. J. Fish. Aq. Sci. 49:783-789.
- Nickelson, T. E., M. F. Solazzi, S. L. Johnson, and J. D. Rodgers. 1993. An approach to determining stream carrying capacity and limiting habitat for coho salmon (*Oncorhynchus kisutch*). Pages 251-260 In L. Berg and P. W. Delaney, Editors. Proceedings of the coho workshop. May 26-28, 1992, British Columbia Department of Fisheries and Oceans, Vancouver, BC.
- Orcutt, D.R., B.R. Pulliam, and A. Arp. 1968. Characteristics of steelhead trout redds in Idaho streams. Transactions of the American Fisheries Society. 97: 42-45.
- OSGC, 1960. Oregon State Game Commission. Statement concerning the fish and wildlife resources of the Deschutes Basin for presentation at the State Water Resources Public

- Oosterhout, G. R. 1999. PasRAS: A stochastic simulation of Chinook and sockeye life histories. Pelton Round Butte Hydroelectric Project FERC no. 2030. Portland General Electric Company, Portland, OR. 101 pages.
- Phillips, R.W., R.L. Lantz, E.W. Claire, and J.R. Moring. 1975. Some effects of gravel mixtures on emergence of coho salmon and steelhead trout fry. Transactions of the American Fisheries Society. 104(3): 461-466.
- Platts, W.S. 1974. Geomorphic and aquatic conditions influencing salmonids and stream classification, with emphasis on ecosystem classification. U.S. Department of Agriculture, Forestry Service, Surface Environment and Mining Program. Boise, Idaho. 199 pp.
- Platts, W.S., and F.E. Partridge. 1978. Rearing of chinook salmon in tributaries of the South Fork Salmon River, Idaho. Intermountain Forest and Range Experiment Station. USDA Forest Service Research Paper Int-205. 11pp.
- Platts, W.S., and F.E. Partridge. 1983. Inventory of salmon, steelhead trout, and bull trout: South Fork Salmon River, Idaho. Intermountain Forest and Range Experiment Station. USDA Forest Service Research Note Int-324. 9 pp.
- Platts, W.S., R.J. Torquemada, M.L. Mchenry, C.K. Graham. 1989. Changes in salmon spawning and rearing habitat from increased delivery of fine sediment to the South Fork Salmon River, Idaho. Transactions of the American Fisheries Society. 118: 274 - 283.
- Rahel, F. J. and W. A. Hubert. 1991. Fish assemblages and habitat gradients in a Rocky Mountain-Great Plains stream: biotic zonation and additive patterns of community change. Transactions of the American Fisheries Society 120:319-332.
- Raleigh, R. F., T. Hickman, R.C. Solomon, and P.C. Nelson. 1986a. Habitat suitability information: rainbow trout. U.S. Fish and Wildlife Service. FWS/OBS - 82/10.60. 64 pp.
- Raleigh, R.F., W.J. Miller, and P.C. Nelson. 1986b. Habitat suitability index models and instream flow suitability curves: Chinook salmon. U.S. Fish and Wildlife Service Biological Report 82 (10.122). 64 pp.
- Reiser, D.W., and T.C. Bjornn. 1979. Habitat requirements of anadromous salmonids. In: Meehan, W.R. Influences of forest and rangeland management on anadromous fish habitat in the western US and Canada. Portland: Pacific Northwest Forest and Range Experiment Station. General Technical Report Pnw-96. 1 - 54.
- Reynolds, K. and S. Peets. 2001. Integrated Assessment and Priorities for Protection and Restoration of Watersheds. Proceedings of the IUFRO 4.11 conference on forest biometry, modeling and information science. 26-29 June 2001, Greenwich, UK. 14 pp. Available at <http://www.institute.redlands.edu/emds/about.htm>

- Sams, R.E. and L.S. Pearson. 1963. Methods for determining spawning flows for anadromous salmonids. Oregon Fish Commission, Portland.
- Scott, W.B. and E.J. Crossman, editors. 1973. Freshwater fishes of Canada. Fisheries Research Board of Canada, Bulletin 184. Ottawa: Bryant Press Limited. 966 pp.
- Seeley, C.M., and G.W. McCammon. 1966. Kokanee. In: Calhoun, A.J., editor. Inland fisheries management. California Department of Fish and Game, Sacramento. P. 274 - 283.
- Seymour, A.H. 1956. Effects of temperature upon young chinook salmon. Ph. D. thesis. University of Washington, Seattle. 127 pp.
- Sheppard, J.D., and J.H. Johnson. 1985. Probability-of-use for depth, velocity, and substrate by subyearling coho salmon and steelhead in Lake Ontario tributary streams. North American Journal of Fisheries Management. 5: 277-282.
- Smith, A.K. 1973. Development and application of spawning velocity and depth criteria for Oregon salmonids. Transactions of the American Fisheries Society 10(2):312 - 316.
- Smith, C.J. 2005. Salmon Habitat Limiting Factors in Washington State. Washington State Conservation Commission. Olympia, Washington. 222pp.
- Steward, C.R., and T.C. Bjornn. 1987. The distribution of chinook salmon juveniles in pools at three discharges. In: Proceedings of the Annual Conference Western Association of the Fish and Wildlife Agencies 67:364-374.
- Stowell, R., A. Espinosa, T.C. Bjornn, W.S. Platts, D.C. Burns, and J.S. Irving. 1983. Guide for predicting salmonid response to sediment yields in Idaho Batholith watersheds. U.S. Forest Service, Northern Region. Missoula, Montana, and Intermountain Region, Ogden, Utah.
- Stuehrenberg, L.C. 1975. The effects of granitic sand on the distribution and abundance of salmonids in Idaho streams. M.S. Thesis, University of Idaho, Moscow. 49 pp.
- Tappel, P.D. and T.C. Bjornn. 1983. A new method of relating size of spawning gravel to salmonid embryo survival. North American Journal of Fisheries Management 3: 123-135.
- Taylor, G.H. and C. Hannan. 1999. The Climate of Oregon: from rainforest to desert. Oregon State University Press, Corvallis, Oregon. 211 pp.
- Thompson, K. 1972. Determining stream flows for fish life. In: Proceedings, instream flow requirement workshop. Vancouver, Washington: Pacific Northwest River Basins Commission. pp. 31-50.
- Thompson K. E. and J. Fortune. 1968. The fish and wildlife resources of the North Coast Basin, Oregon, and their water requirements. Oreg. Game Comm. 102 pp.

- Thorson, T.D., Bryce, S.A., Lammers, D.A., Woods, A.J., Omernik, J.M., Kagan, J., Pater, D.E., and Comstock, J.A., 2003. Ecoregions of Oregon (color poster with map, descriptive text, summary tables, and photographs): Reston, Virginia, U.S. Geological Survey (map scale 1:1,500,000).
- U.S.D.A. Forest Service. 2006. Stream Inventory Handbook. Pacific Northwest Region, Region 6, Version 2.6.
- Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The river continuum concept. *Can. J. Fish. Aquat. Sci.* 37:130-137.
- Vining, T.J., J.S. Blakely, and G.M. Freeman. 1985. An evaluation of the incubation life-phase of chum salmon in the Middle Susitna River, Alaska. Alaska Department of Fish and Game Report 5, Anchorage.
- Vronskii, B.B. 1972. Reproductive biology of the Kamchatka River chinook salmon (*Oncorhynchus tshawytscha*) (Walbaum). *Journal of Ichthyology* 12(2):259-273.
- Wesche, T.A. 1980. The WRRRI trout cover rating method: development and application. Water Resources Research Institute, Laramie, WY. Water Resource Service 78. 46 pp.
- Wilson, W.J. 1984. Pink and chum salmon spawning, egg incubation, and outmigration from two streams on Kodiak Island, Alaska. Pages 26-35 In: K.L. Fresh and S.L. Schroder, rapporteurs. Proceedings of the 1983 Northeast Pacific pink and chum salmon workshop. Washington State Department of Fisheries, Olympia.
- Wolman, M. G. 1954. A method of sampling coarse river-bed material. *Transactions American Geophysical Union*.

APPENDIX A

Life history timing of anadromous salmonids in the Middle Deschutes River basin.

	Life Stage	Jan	Feb	Mar	April	May	June	July	Aug	Sept	Oct	Nov	Dec
Chinook	Immigration						spring						
								summer					
	Spawning												
	Emergence												
	Outmigration				1+								
Steelhead	Immigration			<i>Pelton</i>					fall (summer)			<i>Pelton</i>	
					spring (winter)			<i>Pelton</i>					
	Spawning												
	Emergence												
	Outmigration					1+ and 2+							
							0+						
Sockeye	Immigration												
	Spawning									3 - 7 °C			
	Emergence												
	Outmigration						1+ and 2+						

- a) Shaded boxes represent residence or migration period, darker shaded cells are peaks in migration past Pelton Dam during evaluation period
- b) Immigration is migration into the Deschutes River
- c) Outmigration is at Pelton Dam (rkm 161)
- d) Pelton signifies time at which passing the Pelton weir

APPENDIX B

Chinook salmon habitat requirements and references

Spawning	
<i>Substrate</i>	
Gravel = 3 - 15 cm measured in redd] 15 cm is upper useable limit Gravel = 62% [measured in redd] Cobble = 38% [measured in redd]	Chambers 1956 in Raleigh et al. 1986b
1.3 - 3.8 cm (80%) and up to 10.2 cm (20%) [salmon spawning channel recommendation]	Bell 1986
2 to 10 cm preferred [spawning channel study]	Lucas 1959 in Reiser & Bjornn 1979
6% fines [measured in the redd, Columbia spring chinook] 59 - 86 % gravel 8 - 35% rubble	Burner 1951 in Raleigh et al. 1986b
7.6 - 25.4 cm preference [area prior to spawning, Deschutes chinook]	Huntington 1985
10 cm size limit	Lotspeich & Everest 1981
Salmonids can spawn in gravel w/ median diam \leq 10% of their body length.	Kondolf & Wolman 1993
Avg. dg=24.4 mm, 12.9 % fines reduced to 8.3%	Chambers et al 1954,1955 in Kondolf & Wolman 1993
Reduced fines, <1mm, from 30% to 7.2% [during redd construction] 12 to 26% optimum level of fine sediments in spawning areas	Everest et al. 1987

Spawning (continued)	
<i>Depth (reflect pre-spawning conditions)</i>	
≥ 0.18m (Willamette, n=270)	Sams and Pearson 1963 in Reiser & Bjornn 1979
≥ 0.24m	Thompson and Fortune 1968
≥ 0.2m	Briggs 1953 in Raleigh et al. 1986b
≥ 0.24m spring chinook (Oregon, n=158)	Thompson 1972 in Reiser & Bjornn 1979
≥ 0.24m	Smith 1973
≥ 0.2m at optimum densities	Divinin 1952 in Raleigh et al. 1986b
<i>Temp</i>	
4.4 - 18 °C preferred for spawning	Mattson 1948, Burner 1951 in Raleigh et al. 1986b
low survival (egg + fry) if temp ≥ 16°C no embryo survival at 0°C initially >2 ≤ 3.5 weeks at ≥ 4.5 °C but ≤ 12.8 °C	Seymour 1956 in Raleigh et al. 1986b
10-12 °C favorable range for spawning	Bell 1986
≥ 15 °C may be lethal for embryo	Eddy 1972 in Raleigh et al. 1986b
<i>Flow</i>	Raleigh et al. 1986b
<i>Habitat</i>	
Pool tailouts	Vronskii 1972 in Raleigh et al. 1986b
Pool tailouts	Sullivan et al. 1987
40-60% pools is optimum for spawning and rearing	Raleigh et al. 1986b

Egg survival (incubation)	
<i>surface fines</i>	
≤ 5% silt (≤ 0.8 mm) is optimum ≤ 5% sand (≤ 30.0 mm) is optimum	Raleigh et al. 1986b
< 15% fines (<0.84 mm) is optimal, any greater = decreased survival	McNeil & Ahnell 1964 in Raleigh et al. 1986b
< 5% = high O ₂ permeability > 15% = low O ₂ permeability (<0.84mm)	McNeil & Ahnell 1964 in Bjornn & Reiser 1991, 1979
0 - 30% fines <6.35mm resulted in > 80% survival	Tappel and Bjornn 1983 in Bjornn & Reiser 1991
20% fines <0.83 mm in diameter is upper limit	Everest et al. 1987

Emergence	
<i>Surface fines</i>	
> 25 % fines (< 50% survival)	Bjornn 1969 in Reiser & Bjornn 1979
< 15% (> 75% survival) (\leq 6.4 mm)	
>30-40% sand resulted in nearly no emergence	Bjornn 1968
20% is harmful stage (\leq 0.8mm)	Stowell et al. 1983 in Bjornn & Reiser 1991
utilize 2cm size substrate for cover	Burger et al. 1982 in Raleigh et al. 1986b
20 - 25 % fines (> 75 % survival) (\leq 6.4 mm)	

Summer Rearing 0+ (Fry)	
<i>Substrate Preference</i>	
< 10% fines (< 3mm) in riffle runs	Raleigh et al. 1986b
> 30% fines, low probability of use as cover	
10 - 40 cm substrate \geq 15% of area is adequate cover with < 5% fines	Raleigh et al. 1986b
found over silt to 20cm diameter [0+]	
Habitat is marginal if fines \geq 15% [Pink salmon]	
boulders > 25 cm in riffle runs	Hillman & Griffith 1987
sand and gravel substrate	
as growth occurs, larger substrate	
>~40% fines resulting in embeddedness reduced fish locally (<1fish/m ²)	Bjornn et al 1977 in Bjornn & Reiser 1991
Utilize 2 to 5cm diameter substrate	Bjornn & Reiser 1991
<i>Pool Area</i>	
40 - 60% pool area	Raleigh et al. 1986b
Tendency towards less than 50% for higher densities	Platts 1974
59% chinook found in area with <20% pools	Platts and Partridge 1983
<i>Habitat</i>	
prefer pools	Platts and Partridge 1978
90% used pools & glides	Hillman & Griffith 1987
preferred pools	Murray and Rosenau 1989
all pool habitat esp. alcoves, BW, DP except high gradient	Jonasson et al. 1995-1998
Pools with LWD and willow margins	Johnson et al.1992
prefer pools with > 10 cm depth	Konopacky 1984 in Bjornn & Reiser 1991
Pools and eddies had greatest densities	Everest & Chapman 1972
<i>Temperature</i>	
12-14°C preferred	Brett 1952 in Bjornn & Reiser 1991
12 - 18°C	Raleigh et al. 1986b
slow growth \geq 19.5°C, preferred 9.4 – 13.8°C	Brett et al. 1982
24°C for 1h not harmful	Bjornn 1978 in Bjornn & Reiser 1991
0 to 23-25°C (Salmonids upper and lower lethal limits)	Bjornn & Reiser 1991
<i>Depth</i>	
Enough to cover them	Bjornn & Reiser 1991
Shift to deeper water with growth	Chapman & Bjornn 1969
Correlated with growth	Everest & Chapman 1972

<i>Gradient</i>	
rear in stream reach gradients < 4 - 5%	Lunetta et al. 1997
densities peaked at 4%	Platts 1974
<i>Cover</i>	
Depth : ≥15cm	Everest & Chapman 1972
20 % of all types	Raleigh et al. 1986b
> 15% of 10 - 40 cm sized substrate for cover	Raleigh et al. 1986b
Highest pool complexity had highest densities	Platts 1974
Prefer overhead bank cover (provided 32% cover in trench) to no cover Undercut banks in addition to other cover	Brusven et al. 1986

Overwintering 0+	
<i>Substrate</i>	
< 5% fines optimum, > 30 % tends to prevent use enter gravel or migrate	Raleigh et al. 1986b
enter the substrate	Everest & Chapman 1972
emigrate if lack of substrate cover in cobble/bldrs sand-gravel to silt -cobble (fry size dependent) will not migrate if suitable cobble present	Hillman & Griffith 1987
Overwinter in the substrate	Everest & Chapman 1972
Substrate is major source of cover	Raleigh et al. 1986b
<i>Pool Complexity</i>	
≥ 20% area Class 1 & 2 pools (preferred)	Raleigh et al. 1986b
<i>Habitat</i>	
Pools, glides and RI's, abundant in pools ssoc. with cover	Jonasson et al. 1995-1998
Pools, glides and RI's	Hillman & Griffith 1987
Assoc. with cover overhanging brush + banks	Steward and Bjornn 1987 in Reiser & Bjornn 1979
Assoc. with cover, prefer pools, found in all types LP and Glides	Bjornn & Reiser 1991
<i>Cover</i>	
> 15% cover including 10 -40 cm sized substrate, silt free	Raleigh et al. 1986b
Prefer overhead bank cover (provided 32% cover in trench)	Brusven et al. 1986
Undercut banks with riparian overhanging	Hillman & Griffith 1987
<i>Temperature</i>	
12-14°C preferred	Brett 1952 in Bjornn & Reiser 1991
12 - 18°C	Raleigh et al. 1986b
0°C minimum	Bjornn & Reiser 1991; Raleigh et al. 1986b
≤ 4°C resulting in hiding in substrate	Chapman & Bjornn 1969

Spring 1+ Rearing and Outmigration	
<i>Substrate</i>	
Occupy larger substrate with growth	Hillman & Griffith 1987
prefer rubble	Everest and Chapman 1972
<i>Depth</i>	
≥ 0.6m	Everest and Chapman 1972
40-58cm	Steward & Bjornn 1987 in Bjornn & Reiser 1991
<61 cm	Stuehrenberg 1975 in Bjornn & Reiser 1991
55 - 60 cm	Konopacky 1984 in Bjornn & Reiser 1991
<i>Cover</i>	
1+ assoc. with cover in pools in winter	Steward & Bjornn, unpublished in
vegetation and undercut banks	Bjornn & Reiser 1991
<i>Pool Complexity</i>	Same as steelhead

APPENDIX C

Steelhead trout habitat requirements with references

Spawning	
<i>Substrate</i>	
< 5% fines in redd (optimal)	Raleigh et al. 1986a
Gravel/cobble (1.5 - 10cm) preference	
0.6 - 10.2 cm, criteria for spawning area	Hunter 1973 in Bjornn & Reiser 1979,1991
Favored 1.2 - 10 cm	Orcutt et al. 1968
Pre spawning silt at 14.5% reduced to 7.5 post spawning	Everest et al. 1987
Salmonids can spawn in gravel w/ median diameter \leq 10% of their body length.	Kondolf & Wolman 1993
0.64 - 7.62 cm [probability of use]	Huntington 1985
% Fines [Spawning and rearing]	Platt et al. 1983
5 < 5%	
4 5-25%	
3 25-50%	
2 50-75%	
1 >75%	
<i>Habitat</i>	
Pool tailouts	Greeley 1932 in Raleigh et al. 1986a
Depth (reflects pre-spawning conditions)	
\geq 0.24 m	Smith 1973
Shallowest = 0.21m	Orcutt et al. 1968
<i>Temperature</i>	
10 - 15°C for spawning	Scott and Crossman 1973
\geq 4°C for upstream migration	Hanel 1971 in Raleigh et al. 1986a
3.9 - 9.4 °C preferred for spawning	Bell 1986
7 - 12°C optimum for embryo development	Raleigh et al. 1986a
<4°C and >16°C is low survival (HSI) for embryo	
<i>Flows</i>	adapted from Binns & Eiserman 1979, Wesche 1980 in Raleigh et al. 1986a

Egg Survival	
<i>Substrate</i>	
< 5% fines = high O2 permeability > 15% fines = lower O2 permeability (fines = 0.84mm)	McNeil & Ahnell 1968 in Bjornn & Reiser, 1979 & 1991
0 - 25% fines (>80% survival) (fines<6.35mm) > 40% fines (~ 50% survival)	Tappel and Bjornn 1983 in Bjornn & Reiser 1991
>30-40% fines (1-3mm) resulted in <50% survival [lab]	Hall and Lantz 1968

Emergence	
<i>Substrate</i>	
< 15 % fines (> 90% emergence) >20-25% fines (<50% emergence) (fines < 6.4 mm) resulted in reduced survival + emergence	McCuddin 1977, Bjornn 1969 in Reiser & Bjornn 1979
20% is harmful stage (<6.4 mm)	Stowell et al. 1983 in Bjornn & Reiser 1991
inverse relationship with increased sand	Phillips et al. 1975
> 20% fines (<50% fry emerge)	McCuddin 1977 in Reiser & Bjornn 1979
0-17.5% sand (>80% emerged) >50% sand (<50% emerged)	Bjornn 1968

Summer Rearing 0+	
<i>Substrate</i>	
<10 % fines (interstices and production) 30% fines upper limit	Raleigh et al. 1986a
RI's with boulders > 25cm preferred	Hillman & Griffith 1987
Found over rubble substrate	Everest & Chapman 1972
Closely assoc. with cover (substrate and other)	Fausch 1993
Larger substrate than chinook of same length	Chapman and Bjornn 1969
<i>Depth</i>	
0.09 - 0.15m preference	Sheppard & Johnson 1985
< 0.15m preference	Everest & Chapman 1972
shallower than chinook of same length	Chapman and Bjornn 1969
<i>Pool Area</i>	
40 - 60%	Raleigh et al. 1986a
Tendency towards 50% ratio with riffles	Platts 1974
<i>Cover</i>	
10 - 40 cm substrate in 10% of habitat area (small juveniles) >15% cover including substrate (adequate)	Raleigh et al. 1986a
<10% rating: 0 (worst) 10 to 25% 1 26 to 40 2 41 to 55% 3 >55 4 (best)	Binns & Eiserman 1979 [Trout habitat rating model]
<i>Habitat</i>	
all habitat types	Platts & Partridge 1978
Pools margins, RB's	Hillman & Griffith 1987
RI's, pools, abundant in BW, no preference	Bisson et al. 1988
RI's with LWD, RB, CB	Bisson et al. 1981
Pools, glides, and riffles	Hicks 1990
<i>Temperature</i>	
0-25°C (lower/upper limits)	Lagler 1956, McAfee 1966, Black 1953
optimal 12 - 18°C (rainbow trout)	Raleigh et al. 1986a
10 - 13°C preferred, 0 - 23.9° (lower/upper)	Bell 1986
0 to 23-25°C [Salmonids]	Bjornn & Reiser 1991

Overwintering 0+	
<i>Substrate</i>	
10 - 40 cm substrate which is \geq 10% of total habitat	Raleigh et al. 1986a
Larger substrate shift in winter	Sheppard and Johnson 1985
Rubble, primary cover assoc. with rocks 10 - 25 cm in diameter	Bustard and Narver 1975, cited in Raleigh et al. 1986a
<i>Cover</i>	
\geq 15% including substrate and other undercut banks and cover	Wesche 1980 in Raleigh et al. 1986a
Assoc. with rubble, will emigrate otherwise	Bjornn 1971
Assoc. with cover - rubble primary source	Bustard & Narver 1975
Assoc. with out of channel cover and submerged cover	Bjornn & Reiser 1991
moved to pools and forest canopy in winter (from clear cuts)	Johnson et al 1986
winter cover is important, correlated with substrate	Chapman and Bjornn 1969
<i>Habitat</i>	
Pools low velocity, any habitat with rubble	Bustard & Narver 1975
lower velocity habitat	Sheppard & Johnson 1985
deep pools and abundant cover	Johnson et al. 1986

Summer Rearing 1+	
Substrate	
<10 % fines in riffle (interstices and production)	Raleigh et al. 1986a
found over larger rubble substrate (>40 cm)	Everest & Chapman 1972
occupy larger substrate as they grow	Sheppard & Johnson 1985
Habitat	
Riffles (runs areas)	Raleigh et al. 1986a
found in all habitat types	Platts & Partridge 1983 in Platts et al. 1989
prefer LP and PP, found in all avoided RI, GL, DP, and SC higher velocity and deeper water	Bisson et al. 1988
prefer LP,PP,TP w/ undercut banks and LWD found in RB and CB	Bisson et al. 1981
Depth	
0.6 - 0.75m preferred (I+)	Everest & Chapman 1972
deeper than 0+	Bisson et al. 1988
Cover	
≥ 15% (substrate and other)	Raleigh et al. 1986a
associated with cover	Fausch 1993
assoc. with cover	Bisson et al. 1988
assoc. with cover undercut banks and LWD	Bisson et al. 1981
Overwintering 1+ and Outmigration	
Substrate	
enter substrate - boulders and under logs	Bustard & Narver 1975
10 - 40 cm substrate which is ≥ 10% of total habitat, silt-free	Raleigh et al. 1986a
Class 1 pools	Lewis 1969 in Raleigh et al. 1986a
Rubble (15-45cm diameter) substrate [trough] Rubble or undercut banks [nature]	Bjornn 1971
Depth	
> 45cm	Bustard & Narver 1975
> 45 cm (otherwise, lower densities) 0.6 - 0.75 m preferred (I+)	Everest & Chapman 1972
Cover	
prefer > 40cm boulders	Everest & Chapman 1972
≥ 15% (substrate and other)	Raleigh et al. 1986a
highest density associated with pool depth undercut banks, large rock and brush as strong affinity to large rock as PD,UB and LR combined	Bjornn and Steward, unpublished in Bjornn & Reiser 1991
moved to pools and forest canopy in winter (from clear cut)	
Deep pools with LWD in streams (w/o >40cm rubble), and rubble in rivers	Bustard & Narver 1975

Pool Complexity

Hartman 1965, Lister and Genoe 1970, Everest and Chapman 1972, Edmundson et al 1968 in Raleigh et al. 1986a.

Platts 1974

Platts and Partridge 1983

Rating	Length or Width	Depth	Cover
1	> ACW	≥0.61 m	abundant
	< ACW	≥ 0.91m	absent
2	>ACW	≥0.61 m	Abundant
	>ACW	≥0.61 m	Intermediate
	>ACW	≥0.61 m	absent
3	= ACW	≥0.61 m	Abundant
	= ACW	≥0.61 m	intermediate
4	= ACW	~ equal to average stream depth	absent
	<ACW	~ equal to average stream depth	abundant
		~ equal to average stream depth	intermediate
		≥0.61 m	Intermediate
		≥0.61 m	abundant
5	< ACW	~ equal to average stream depth	absent

Source: Platts 1974: Pool quality rating

Cover: woody debris, boulders, vegetation (in channel or overhanging), and undercut banks.

Rating	Diameter	Depth	Cover
5	> average stream width	> 0.92m	Absent
		> 0.6m	Abundant
4	> average stream width	< 0.6m	absent
		0.6 to 0.91m	Absent
3	< average stream width	> 0.6m	Intermediate to abundant
2	< average stream width	< 0.6m	Intermediate to abundant
1	< average stream width	< 0.6m	absent

Source: Platts and Partridge 1983: Pool classification

Rating	Width	Depth	Cover
First class	≤ 5.0m	≥ 1.5m	30%
	> 5.0m	> 2.0m	
Second class	Moderate	Moderate	5 – 30%
Third class	Small	Shallow	< 5%

Source: Raleigh et al. 1986a: Pool classification

APPENDIX D

Sockeye salmon habitat requirements references

Spawning, egg survival, emergence	
<i>Substrate</i>	
salmonids can spawn in gravel w/ median diam \leq 10% of their body length.	Kondolf 1993
< 5% fines in redd > 15% lower O2 permeability	McNeil & Ahnell 1964 in Bjornn & Reiser 1991
1.3-10.2cm	Bell 1986
medium to small gravel with no silt	Eiler 1992
<15% fines (<2mm) (PU) Typically spawning where there is upwelling, so substrate is highly variable	Lorenze and Eiler 1989
20% is harmful stage	Stowell et al. 1983; Bjornn & Reiser 1991
<i>Habitat</i>	
Areas of upwelling or subsurface flow preferred for spawning	Lister et al 1970, Wilson 1984, Vining et al 1985 in Bjornn & Reiser 1991
small streams of lakes, gravel shores with upwelling or tributaries of lake outlet	Meehan and Bjornn 1991
Lake shore or tributary riffle areas preferred Concentrate in areas of upwelling	Groot 1991
<i>Depth</i>	
enough to cover the fish (minimum)	Groot 1991
\geq 0.15m [estimated]	Bjornn & Reiser 1979, 1991
<i>Temperature</i>	
10.6 - 12.2°C preferred 4.4 - 13.3°C for incubation	Bell 1986
15.5°C mortalities ensue 5.5-12.8°C preferred for spawning	Seeley & McCammon 1966

Summer Rearing 0+ and migration to lake	
<i>Cover</i>	
use undercut banks, overhanging vegetation, and gravel	Hartman et al. 1962
Use gravel or above gravel when not migrating [trough]	McDonald 1960
<i>Habitat</i>	
0+ rear in lakes, rivers, estuaries, and ocean	Groot 1991
0+ rear in lakes, rivers, estuaries and ocean usually in lakes	Meechan & Bjornn 1991
<i>Temperature</i>	
11.1 - 14.4°C preferred	Bell 1986
12 - 14°C preferred, 3.1 - 25.8°C (limits)	Brett 1952 in Bjornn & Reiser 1991
0 to 23-25°C (salmonids)	Bjornn & Reiser 1991

APPENDIX E

Chinook salmon habitat criteria.

Spawning, Egg Survival, and Emergence

prior to redd construction

	3	2	1
Substrate			
Fines (< 2mm)	≤ 10 %	10 - 20 %	>20 %
Gravel (2 – 64mm)	≥ 30 %	15 - 30 %	<15 %
Cobble (64-256mm)	20 - 40 %	10-20,40-70 %	< 10 % , > 70 %
Habitat (Pool Tailouts)	40 - 60 % pools	20 - 40 %	< 20 % , > 60%
Residual Pool Depth	≥ 0.2m		dry
Gradient	< 4 %		≥ 4 %
Temperature	6 - 14°C	4 - 6°C, 14-16°C	< 4°C, > 16°C
Flow	50-100 % base flow	25-50% base flow	< 25 % base flow > annual base flow

* lethal levels extending longer than 1 hour in 24 hour period

Summer Rearing 0+

	3	2	1
Substrate			
Fines (<i>interstices and productivity</i>)	≤ 10 %	10 - 30 %	> 30 %
Gravel (<i>cover</i>)	≥ 15 %	5 - 15 %	< 5 %
Cobble and Boulder (<i>cover</i>)	≥ 15 %	8 - 15 %	< 8 %
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Pool Complexity	3	2	1
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 –15	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m (<i>cobble and boulder from above</i>)	≥ 20	5 – 20	< 5
Habitat (Gradient)	Prefer pools, (≤ 4%)		> Rapids (> 4%)
Temperature	9.5 - 14°C	4 – 9.5° , > 14°C	Lethal levels* (24°C)
Flow	50 - 100 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Overwintering 0+

	3	2	1
Substrate			
Fines (<i>interstices</i>)	≤ 10 %	10 - 30 %	> 30 %
Cobble and Boulder (<i>cover</i>)	≥ 15 %	8 - 15 %	< 8 %
Pool Complexity	3	2	1
Habitat (Gradient)	Pools, GL, RI assoc. with cover (< 4%)		≥ Rapids (≥ 4%)
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 –15	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m	≥ 20	5 – 20	< 5
Flow	100 - 50% base flow	25 -50% base flow	< 25 % base flow

Spring 1+ and Emigration

	3	2	1
Substrate			
Fines (<i>interstices</i>)	≤ 10 %	10 - 30 %	> 30 %
Cobble and Boulder (<i>cover</i>)	≥ 20 %	10 - 20 %	< 10 %
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Pool Complexity	3	2	1
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 –15	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m	≥ 20	5 – 20	< 5
Habitat (Gradient)	Prefer Poor gradient (≤ 4%)		> Rapids (> 4%)
Temperature	9.5 - 14°C	4 – 9.5° , > 14°C	Lethal levels* (24°C)
Flow	100 - 50 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Pool Complexity

3	Deep with considerable cover
	Depth > 0.6 m (≤ 10m wetted width stream)
	Depth > 1 m (> 10m wetted width stream)
	Criteria Conditions*:
	Keypieces of LWD > 0.6 or Pieces of LWD ≥ 2.0
	Undercut bank > 20 %
	Boulders in pools > 15 %
2	Moderate depth and cover
	Depth ≥ 0.6 m (≤ 10m wetted width stream)
	Depth ≥ 0.6 – 1.0 m (> 10m wetted width stream)
	Criteria Conditions*:
	LWD present
	Undercut banks = 5 - 20 %
	Boulders = 8 - 15 %
1	Shallow and lacking cover
	Depth < 0.6 m (≤ 10m wetted width stream)
	Depth < 0.6 m (> 10m wetted width stream)
	Criteria Conditions*:
	No LWD
	Undercut banks < 5 %
	Boulders < 8 %

APPENDIX F

Steelhead trout habitat criteria.

Spawning, egg survival, emergence

prior to redd construction

	3	2	1
Substrate			
Fines	≤ 10 %	10 - 20 %	> 20%
Gravel	≥ 30 %	15 - 30 %	< 15 %
Cobble	10 - 30 %	30 - 60 %	< 10 % , > 60 %
Habitat (Pool Tailouts)	40 - 60 %	20 - 40 %	< 20 % , > 60%
Residual Pool Depth	≥ 0.2 m		No Pools
Temperature	6 - 12.5°C	4- 6°C, 12.5-16°C	< 4°C, > 16°C
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow > annual base flow

Summer Rearing 0+

	3	2	1
Substrate			
Fines (<i>interstices and productivity</i>)	≤ 10 %	10 - 30 %	> 30%
Cobble and Bldr (<i>cover</i>)	≥ 20 %	10 - 20 %	< 10 %
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 - 15	< 10
LWD / 100m	≥ 20	10 - 20	< 10
Boulders / 100m	≥ 20	5 - 20	< 5
Temperature	10 - 13°C	< 10, >13°C	Lethal levels* (24°C)
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Overwintering 0+

	3	2	1
Substrate			
Fines (<i>interstices</i>)	≤ 10 %	10 - 30 %	> 30%
Cobble and Bldr (<i>cover</i>)	≥ 20 %	10 -20%	< 10%
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 –15	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m	≥ 20	5 – 20	< 5
Pool Complexity	3	2	1
Habitat (Gradient)	Pools & RI with cover (< 4%)	all else (≥ 4 %)	
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Summer Rearing 1+

	3	2	1
Substrate			
Fines (<i>interstices & productivity</i>)	≤ 10 %	10 - 30 %	> 30%
Cobble and Boulder (<i>cover</i>)	≥ 20 %	10 - 20 %	< 10%
Depth (<i>in riffles</i>)	≥ 0.45 m		< 0.45 m
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 15	10 –15	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m	≥ 20	5 – 20	< 5
Temperature	10 - 13°C	< 10, >13°C	Lethal levels* (24°C)
Flows	100- 50 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Overwintering 1+ and Emigration

	3	2	1
Substrate			
Fines (<i>interstices</i>)	≤ 10 %	10 - 30 %	> 30%
Cobble and Boulder(<i>cover</i>)	≥ 25 %	10 - 25%	< 10%
Pool Area	40 - 60 %	20 - 40 %	< 20 % , > 60%
Additional Cover (<i>at least one true</i>)			
% Undercut	≥ 20	10 – 20	< 10
LWD / 100m	≥ 20	10 – 20	< 10
Boulders / 100m	≥ 20	5 – 20	< 5
Pool Complexity	3	2	1
Temperature	10 - 13°C	< 10, >13°C	Lethal levels* (0°C)
Smoltification	> 4°C, < 13°C		> 13°C
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow

* lethal levels extending longer than 1 hour in 24 hour period

Pool complexity - refer to chinook criteria

APPENDIX G

Sockeye salmon habitat criteria.

Spawning, egg survival, fry emergence

prior to redd construction

	3	2	1
Substrate			
Fines	≤ 10 %	10 - 30 %	> 30%
Gravel	≥ 30 %	15 - 30 %	< 15 %
Cobble	10 - 40 %	40 - 60 %	< 10 %, > 60 %
Habitat (gradient)	lakeshore or trib with upwelling		high gradient
Residual Pool Depth	≥ 0.15m		≤ 0.15m
Temperature	4.4 - 13.3 °C	< 4.4°C, > 13.3°C	< 1°C, > 20°C
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow, > annual base flow

* lethal levels extending longer than 1 hour in 24 hour period

Summer Rearing 0+ including migration to lake habitat

	3	2	1
Depth			no passage
Cover - undercut banks	≥ 30%	10 - 30%	≤ 10%
Habitat	Lakes		
Temperature	12 - 14°C	< 12, >14°C	Lethal levels* (25°C)
Flows	100 - 50 % base flow	25-50% base flow	< 25 % base flow, > annual base flow

* lethal levels extending longer than 1 hour in 24 hour period



3406 Cherry Ave. NE

Salem, Oregon 97303